

What e^+e^- collider could make a “no-lose” search
for MSSM Higgs bosons?

V. Barger*, Kingman Cheung*, R. J. N. Phillips† and A. L. Stange*

**Physics Department, University of Wisconsin, Madison, WI 53706, USA*

†Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX England

ABSTRACT

The lightest CP-even Higgs boson h in the minimum supersymmetric standard model (MSSM) has a mass upper bound depending on the top quark and squark masses. An e^+e^- collider with enough energy and luminosity to produce $h + Z$ at measurable rates up to the maximum h mass would cover the entire MSSM parameter space, if $h + A$ production was also searched for. We explore the energy and/or luminosity needed for various top quark and squark masses. For $m_t = 150$ GeV and 1 TeV SUSY mass scale, a 230 GeV collider with 10 fb^{-1} luminosity would suffice.

The theoretical appeal of supersymmetry (SUSY) is that it solves the problem of large radiative corrections in the scalar sector, associated with the grand unification scale. The minimal supersymmetric extension of the Standard Model (MSSM) [1] has five Higgs bosons, one of which (h) is necessarily relatively light; their discovery could contribute the first direct evidence both for SUSY and for the Higgs mechanism. These Higgs bosons are therefore the object of intense experimental investigation; a lower limit $m_h \gtrsim 40$ GeV has already been set by e^+e^- experiments at LEP I [2–5] and the range of search will be extended at LEP II with CM energy $\sqrt{s} = 190$ GeV to 240 GeV possible [6]. In this letter we address the question: what is the lowest energy e^+e^- collider that could completely cover the MSSM parameter space [7–11] and thereby independently guarantee discovery or rejection of the MSSM? This question is relevant because LEP I, LEP II, SSC and LHC will not fully cover all MSSM parameters [7,10,11], and the possibilities of higher energy e^+e^- linear colliders are being examined [12].

The Higgs sector of the minimum supersymmetric standard model (MSSM) has three neutral and two charged Higgs bosons, h , H , A , $H^\pm$ of which h and H are CP-even and $m_h < m_H$; a mixing angle α appears in the h and H couplings. At tree level all their masses and couplings are controlled by two parameters, that may be taken to be m_A and the ratio $\tan\beta = v_2/v_1$ of vacuum expectation values giving masses to up-type quarks (v_2) and down-type quarks (v_1) respectively; renormalization group arguments in no-scale models [13] suggest that $1 < \tan\beta < m_t/m_b$ but m_A is unconstrained. At one-loop level, however, there are significant radiative corrections [14], that depend on several other parameters but especially on the top quark and squark masses; as a result the h mass has an upper bound

$$m_h^2 \lesssim M_Z^2 \cos^2 \beta + \frac{6G_F}{\pi^2\sqrt{2}} m_t^4 \ln \left(\frac{\tilde{m}}{m_t} \right) , \quad (1)$$

in an approximation where the usual SUSY parameters A_t , A_b and μ are set to zero and $\tilde{m}$ is the common SUSY mass scale. The masses of H , A and $H^\pm$ have no upper bounds. In our present discussion we shall use non-zero values of all SUSY parameters, following Ref. [8], with $\tilde{m} \simeq 1$ TeV and $1 < \tan\beta < 30$. The important parameter is the shift in the m_h upper

bound; we note that large changes in $\tilde{m}$ are effectively equivalent here to small changes in m_t .

At e^+e^- colliders the signals for Higgs bosons are relatively clean and the opportunities for discovery and detailed study will be excellent. The principal production channels are

$$e^+e^- \rightarrow Z \rightarrow Zh, ZH, Ah, AH, \quad (2a)$$

$$e^+e^- \rightarrow \nu\bar{\nu}W^*W^* \rightarrow \nu\bar{\nu}h, \nu\bar{\nu}H, \quad (2b)$$

$$e^+e^- \rightarrow t\bar{t}h, t\bar{t}H, t\bar{t}A, \quad (2c)$$

$$e^+e^- \rightarrow Z, \gamma \rightarrow H^+H^-, \quad (2d)$$

$$e^+e^- \rightarrow t\bar{t} \rightarrow b\bar{b}H^+(W^+)H^-(W^-). \quad (2e)$$

The s -channel processes (2a) offer the biggest contributions at the lower energies. In the limit $m_A \rightarrow \infty$ we have $m_H \simeq m_{H^+} \simeq m_A$ while m_h approaches the upper bound of Eq. (1) and only h can be produced at any given collider. Thus the channel $e^+e^- \rightarrow Zh$ and its kinematical limits are critical in any complete search of MSSM parameter space. The Zh production cross section contains an overall factor $\sin^2(\beta - \alpha)$ which suppresses it in certain parameter regions (with $m_A < 100$ GeV and $\tan\beta$ large); fortunately the Ah production cross section contains the complementary factor $\cos^2(\beta - \alpha)$. Hence the Zh and Ah channels together are well suited to cover all regions in the $(m_A, \tan\beta)$ plane, provided that the CM energy is high enough for Zh to be produced through the whole m_h mass range, and that an adequate event rate can be achieved. These conditions are already shown to be fulfilled [12] for $\sqrt{s} = 500$ GeV with assumed luminosity 10 fb^{-1} . In the present work we study how well these conditions can be fulfilled at lower energies with various luminosities.

Our discussion centers on the s -channel production channels $e^+e^- \rightarrow Zh, Ah, ZH, AH$, neglecting all others for simplicity (although other channels would obviously contribute to an eventual search and analysis). We also consider only the decays $Zh, Ah, ZH, AH \rightarrow \tau\tau jj$ (where j denotes a b -jet); this generally has substantial branching fractions at least in the Zh and Ah cases. We rely here on the possibility of recognizing and kinematically reconstructing τ jets experimentally [2–5,12]; no b -tagging of the other jets is assumed. This approach is

conservative, since it implicitly ignores substantial $Z \rightarrow \ell^+ \ell^-, \nu \bar{\nu}$ decay modes that could enhance the detectability of Zh and ZH production. Since the $\ell^+ \ell^- jj$ and $\nu \bar{\nu} jj$ channels have smaller signal/background ratios than the $\tau \tau jj$ channel, the net significance of Zh or ZH signals would not be dramatically increased by including the former channels.

For any given energy and MSSM parameters, we calculate the Zh , Ah , ZH and AH production cross sections and decay branching fractions from standard formulas [1] with one-loop corrections as described in Ref. [8]. We omit bremsstrahlung and beamstrahlung corrections, that are not very large in currently favored collider designs. It is assumed that the Higgs bosons do not decay to light SUSY particles. The signals take the form of peaks in the distributions of invariant mass $m(\tau\tau)$ and $m(jj)$, centered at values m_h , m_H and m_A , with an associated peak at M_Z also. These two $m(\tau\tau)$ and $m(jj)$ distributions are added to enhance the statistics, thus giving two counts per event. An irreducible background from $e^+e^- \rightarrow ZZ$ production and decay has a peak centered at M_Z ; all other backgrounds can however be suppressed by suitable cuts at little cost to the signals [12]. For present purposes we estimate the background from the numerical simulation of Ref. [12], scaling the number of events according to the assumed luminosity and the energy-dependence of the $e^+e^- \rightarrow ZZ \rightarrow \tau\tau jj$ cross section with $|\cos\theta| < 0.9$ for the τ 's and jets. We assume that the acceptances of the Zh (ZH) and Ah (AH) signals remain 46% and 52%, respectively, as in Ref. [12] and that the Higgs boson peaks have the same mass resolution as the Z peak. This approach is approximate, but avoids lengthy Monte Carlo simulations for each of the many different energies and parameter settings that we have to consider.

For each input set of SUSY parameters, CM energy $\sqrt{s}$ and integrated luminosity $\mathcal{L}$, we define the signals, backgrounds and discovery criteria of the MSSM mass-peaks as follows. For an isolated peak, the signal strength S is taken to be the expected number of signal counts falling in a 10 GeV mass bin centered at the corresponding Higgs boson mass. When two Higgs peaks approach within 10 GeV we combine them; the signal strength S is then the total number of counts expected in a 10 GeV bin centered at the weighted mean mass. The background strength B is taken to be the total number of Z -decay counts (both from ZZ

and from Zh, ZH production with the resolution of Ref. [12]) falling in the same mass bin. If the signal bin center is separated by more than 5 GeV from M_Z , our discovery criteria are $S/\sqrt{B} > 4$ with $S > 4$ counts. With such a separation, we expect that a distinct peak will be seen or that a recognizable distortion of the Z peak will be evident. But if the separation from M_Z is less than 5 GeV, we can only infer the presence of a new signal if the height of the supposed Z peak differs substantially from the expected ZZ background contribution. In this latter case we rely entirely on normalization and therefore require a higher degree of significance. Here the signal S is defined to be the sum of the MSSM (h, H, A and Z) contributions falling in a 10 GeV bin centered at M_Z , and B is the expected ZZ background in the same bin; in this case we define a discoverable signal to have $S/\sqrt{B} > 6$ with $S > 5$ counts.

In principle, b -tagging of the quark jets offers another way to distinguish the presence of a Higgs boson contribution hiding under the Z peak in the $\tau\tau jj$ channel. With typical LEP-type microvertex tagging, it should be possible to achieve say 38% tagging efficiency for $Higgs \rightarrow b\bar{b}$ pairs coupled with 11% tagging efficiency for $Z \rightarrow jj$ pairs. If we consider just the $m(jj)$ distribution for simplicity, a tagged signal would then have significance increased by approximately the factor $0.38/\sqrt{0.11} = 1.14$ above that of the corresponding untagged signal (discussion of the $m(\tau\tau)$ distribution is more complicated). Greater enhancement may well be attainable in the future. It may also be possible to establish a signal in the ratio of tagged/untagged events, but this seems to require much higher statistics to achieve serious significance. Thus b -tagging appears to offer real but not dramatic improvements in sensitivity in the future; conservatively we neglect it here.

For full coverage of the $(m_A, \tan\beta)$ plane, the CM energy should be about 10 GeV or more above the maximum Zh threshold,

$$\sqrt{s}(\text{threshold}) = m_h(\text{max}) + M_Z, \quad (3)$$

where $m_h(\text{max})$ is the largest value of m_h in Eq. (1). For $\tilde{m} = 1$ TeV this threshold is 207 GeV for $m_t = 150$ GeV and 240 GeV for $m_t = 200$ GeV (the highest value of m_t allowed

by analyses of radiative corrections [15]). Apart from these threshold considerations, the principal factors that determine the discovery regions (where one or more MSSM signals are detectable) in the $(m_A, \tan\beta)$ plane are luminosity $\mathcal{L}$, top quark mass m_t and CM energy $\sqrt{s}$. Figure 1 illustrate the effects of these factors separately, by means of four examples.

- (i) Luminosity: in Figs. 1(a),(b) we hold $m_t = 150$ GeV and $\sqrt{s} = 215$ GeV fixed (with all SUSY parameters fixed as in Ref. [8]), and compare discovery limits for $\mathcal{L} = 1$ fb $^{-1}$ and $\mathcal{L} = 10$ fb $^{-1}$, corresponding respectively to one month and one year running at $L = 10^{33}$ /cm 2 /s. Here the $(m_A, \tan\beta)$ plane is fully accessible kinematically, but good luminosity is still needed to guarantee discovery; in fact $\mathcal{L} \gtrsim 20$ fb $^{-1}$ would give full coverage.
- (ii) Top quark mass: in Figs. 1(b),(c) we hold $\mathcal{L} = 10$ fb $^{-1}$ and $\sqrt{s} = 215$ GeV fixed and compare the discovery limits for $m_t = 150$ GeV and $m_t = 200$ GeV. Coverage becomes easier as m_t decreases; there would be complete coverage in this case with $m_t \lesssim 120$ GeV.
- (iii) CM energy: in Figs. 1(c),(d) we hold $\mathcal{L} = 10$ fb $^{-1}$ and $m_t = 200$ GeV fixed and compare the discovery limits at $\sqrt{s} = 215$ and 270 GeV. We see that increasing s in this range generally widens the accessible region, although this is not uniformly true since the signals have different energy dependences in different parts of the plot. In fact, with our discovery criteria it appears that complete coverage is not achieved at any energy with this particular choice of $\mathcal{L}$ and m_t . We remark in passing that the small area lower left, inaccessible in Figs. 1(a) and 1(d), is not well served by $\tau\tau jj$ signals, since $h \rightarrow AA$ dominates the h decays here; however, this region is already excluded by LEP I data [2–5].

The final question is, what combinations of collider parameters $\sqrt{s}$ and $\mathcal{L}$ would just achieve complete coverage of the $(m_A, \tan\beta)$ plane for given m_t ? Figure 2 shows the limiting curves in the $(\sqrt{s}, \mathcal{L})$ plane, for various values of m_t ; we recall that changes in $\tilde{m}$ can

be effectively absorbed into m_t , and that LEP I searches have already excluded small m_A values [2–5]. Pairs of values $(\sqrt{s}, \mathcal{L})$ that lie above the limiting curves have “no-lose” discovery potential in the MSSM, according to our approximations. For example, with $m_t = 150$ GeV and a 1 TeV SUSY mass scale, a 230 GeV e^+e^- collider with 10 fb^{-1} luminosity would suffice.

We thank Peter Norton for conversations about experimental issues and Paul Dauncey for information about b -tagging efficiencies. This work is supported in part by the U.S. Department of Energy under contract No. DE-AC02-76ER00881, in part by the Texas National Research Laboratory Commission under grant No. RGFY9173, and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

REFERENCES

- [1] For a recent reviews see J. Gunion, H. Haber, G. Kane and S. Dawson, *The Higgs Hunter's Guide*, (Addison-Wesley, 1990); X. Tata in *The Standard Model and Beyond*, edited by J. E. Kim (World Scientific, 1991) p. 379.
- [2] ALEPH Collaboration, D. Decamp *et al.*, Phys. Lett. **B265**, 475 (1991), and CERN-PPE/91-149 to be published in Physics Reports.
- [3] DELPHI Collaboration: P. Abreu *et al.*, Phys. Lett. **B245**, 276 (1990).
- [4] L3 Collaboration: B. Adeva *et al.*, Phys. Lett. **B251**, 311 (1990).
- [5] OPAL Collaboration: M. Z. Akrawy *et al.*, Z. Phys. **C49**, 1 (1991).
- [6] C. Rubbia, in *Proc. LEP-HEP 91*, ed. by S. Hegarty *et al.*, World Scientific (1991), p. 439.
- [7] Z. Kunszt and F. Zwirner, in *Proceedings of the Large Hadron Collider Workshop at Aachen*, CERN 90-10; revised results in CERN-TH.6150/91.
- [8] V. Barger, M. S. Berger, A. L. Stange and R. J. N. Phillips, Madison report MAD/PH/680 (1991), Phys. Rev. D (in press).
- [9] H. Baer, M. Bisset, C. Kao and X. Tata, Florida State Univ. report FSU-HEP-911104 (1991).
- [10] J. F. Gunion, R. Bork, H. E. Haber and A. Seiden, U. C. Davis report UCD-91-29; J. F. Gunion and L. H. Orr, report UCD-91-15; J. F. Gunion, H. E. Haber, and C. Kao, report UCD-91-32.
- [11] V. Barger, K. Cheung, R. J. N. Phillips, and A. L. Stange, *Collider discovery limits for supersymmetric Higgs bosons*, UW-Madison report MAD/PH/696.
- [12] P. Janot, Orsay preprint LAL 91-61 (1991); J.-F. Grivaz, Orsay preprint LAL 91-63 (1991).

- [13] See for example F. Zwirner, CERN-TH.6357/91.
- [14] S. P. Li and M. Sher, Phys. Lett. **B140**, 339 (1984); J. F. Gunion and A. Turski, Phys. Rev. **D39**, 2701, **D40**, 2325, 2333 (1989); M. S. Berger, Phys. Rev. **D41**, 225 (1990); Y. Okada *et al.*, Prog. Theor. Phys. **85**, 1 (1991); Phys. Lett. **B262**, 54 (1991); H. Haber and R. Hempfling, Phys. Rev. Lett. **66**, 1815 (1991); J. Ellis *et al.*, Phys. Lett. **B257**, 83 (1991); R. Barbieri *et al.*, Phys. Lett. **B258**, 167 (1991); J. Lopez and D. V. Nanopoulos, Phys. Lett. **B266**, 397 (1991); A. Yamada, Phys. Lett. **B263**, 233 (1991); M. Drees and M. N. Nojiri, KEK-TH-305 (1991); R. Hempfling, SCIPP-91/39; M. A. Diaz and H. E. Haber, SCIPP-91/14 (1991); D. M. Pierce, A. Papadopoulos and S. Johnson, LBL-31416 (1991); P. H. Chankowski, S. Pokorski and J. Rosiek, Phys. Lett. **B274**, 191 (1992); J. R. Espinosa and M. Quiros, Phys. Lett. **B279**, 92 (1992).
- [15] P. Langacker, in the *Review of Particle Properties*, Phys. Rev. **D45**, VII.159 (1992).

FIGURES

FIG. 1. Discovery limits in the $(m_A, \tan \beta)$ plane for $e^+e^- \rightarrow Zh, Ah, ZH, AH$ signals in the $\tau\tau jj$ channel, for various illustrative cases: (a) $\sqrt{s} = 215 \text{ GeV}$, $m_t = 150 \text{ GeV}$, $\mathcal{L} = 1 \text{ fb}^{-1}$;
(b) $\sqrt{s} = 215 \text{ GeV}$, $m_t = 150 \text{ GeV}$, $\mathcal{L} = 10 \text{ fb}^{-1}$;
(c) $\sqrt{s} = 215 \text{ GeV}$, $m_t = 200 \text{ GeV}$, $\mathcal{L} = 10 \text{ fb}^{-1}$;
(d) $\sqrt{s} = 270 \text{ GeV}$, $m_t = 200 \text{ GeV}$, $\mathcal{L} = 10 \text{ fb}^{-1}$.

FIG. 2. Conditions for covering the whole MSSM $(m_A, \tan \beta)$ plane with $m_A \leq 1 \text{ TeV}$ and $1 < \tan \beta < 30$. Limiting curves are shown in the $(\sqrt{s}, \mathcal{L})$ plane, for various values of m_t . The region that does not give complete coverage for $m_t = 150 \text{ GeV}$ is shaded.